


---

### Infants' Acetaminophen Concentrated Drops

The enclosed dropper is designed to provide 0.4 mL (40 mg) increments




- Remind parents and caregivers:**
- Infants' Acetaminophen Concentrated Drops are 3x more concentrated than Acetaminophen Oral Suspension
  - Always read and follow the label

---

### Infants' Acetaminophen Oral Suspension

The enclosed syringe is designed to provide 1.25 mL (40 mg) increments




- Keep medicines out of the reach of children
- Never give adult medicines to children
- Use only the dosing device that comes with the product

---

### Infants' Acetaminophen Concentrated Drops

The enclosed dropper is designed to provide 0.4 mL (40 mg) increments




- Remind parents and caregivers:**
- Infants' Acetaminophen Concentrated Drops are 3x more concentrated than Acetaminophen Oral Suspension
  - Always read and follow the label

---

### Infants' Acetaminophen Oral Suspension

The enclosed syringe is designed to provide 1.25 mL (40 mg) increments




- Keep medicines out of the reach of children
- Never give adult medicines to children
- Use only the dosing device that comes with the product

---

### Infants' Acetaminophen Concentrated Drops

The enclosed dropper is designed to provide 0.4 mL (40 mg) increments



- Remind parents and caregivers:**
- Infants' Acetaminophen Concentrated Drops are 3x more concentrated than Acetaminophen Oral Suspension
  - Always read and follow the label

---

### Infants' Acetaminophen Oral Suspension

The enclosed syringe is designed to provide 1.25 mL (40 mg) increments




- Keep medicines out of the reach of children
- Never give adult medicines to children
- Use only the dosing device that comes with the product

---

### Infants' Acetaminophen Concentrated Drops

The enclosed dropper is designed to provide 0.4 mL (40 mg) increments



- Remind parents and caregivers:**
- Infants' Acetaminophen Concentrated Drops are 3x more concentrated than Acetaminophen Oral Suspension
  - Always read and follow the label

---

### Infants' Acetaminophen Oral Suspension

The enclosed syringe is designed to provide 1.25 mL (40 mg) increments



- Keep medicines out of the reach of children
- Never give adult medicines to children
- Use only the dosing device that comes with the product

**TYLENOL (ACETAMINOPHEN)**

		INFANT CONCENTRATED DROPS	INFANTS SUSPENSION	CHILDRENS SUSPENSION	CHILDREN'S CHEWABLES	JUNIOR STRENGTH CHEWABLES	REGULAR STRENGTH ADULT	EXTRA STRENGTH ADULT	Acetaminophen can be dosed up to every 4 hours	For better fever control, alternate dosing of acetaminophen and ibuprofen every 4 hours. Do not give more than the recommended amount based on your child's weight or age. Do not give more than 6 doses in a 24 hour period. For infants under 2 months of age, do not give unless directed by your doctor. If your child has a fever lasting more than 3 days, contact your doctor's office.
	STRENGTH	80 mg/0.8 mL	160 mg/5 mL	160 mg/5 mL	80 mg each	160 mg each	325 mg each	500 mg each		
	DOSE	DROPPER OR SYRINGE	SYRINGE	TEASPOON OR SYRINGE	TABLET	TABLET	TABLET OR CAPSULE	TABLET OR CAPSULE		
WEIGHT (lbs)	AGE									
6-11	0-3 months	0.4 mL	1.25 mL	1/4 tsp or 1.25 mL						
12-17	4-11 months	0.8 mL	2.5 mL	1/2 tsp or 2.5 mL						
18-23	12-23 months	1.2 mL	3.75 mL	3/4 tsp or 3.75 mL						
24-35	2-3 years	1.6 mL	5 mL	1 tsp or 5 mL	2 tablets	2 tablets	1 each			
36-47	4-5 years			1 1/2 tsp or 7.5 mL	3 tablets	3 tablets	1 1/2 each			
48-59	6-8 years			2 tsp or 10 mL	4 tablets	4 tablets	2 each			
60-71	9-10 years			2 1/2 tsp or 12.5 mL	5 tablets	5 tablets	2 1/2 each	1 each		
72-95	11 years			3 tsp or 15 mL	6 tablets	6 tablets	3 each	1 each		
over 96	12 years & older						4 tablets	2 each		

**MOTRIN OR ADVIL (IBUPROFEN)**

		INFANT CONCENTRATED DROPS	CHILDRENS SUSPENSION	CHILDREN'S CHEWABLES	JUNIOR STRENGTH CHEWABLES	JUNIOR STRENGTH PILLS	REGULAR STRENGTH ADULT	Ibuprofen can be dosed every 6-8 hours  Do not give ibuprofen if your child has been vomiting or has a stomach ache as it may worsen their stomach problem	For better fever control, alternate dosing of acetaminophen and ibuprofen every 4 hours. Do not give more than the recommended amount based on your child's weight or age. Do not give more than 6 doses in a 24 hour period. For infants under 2 months of age, do not give unless directed by your doctor. If your child has a fever lasting more than 3 days, contact your doctor's office.
	STRENGTH	50 mg/1.25 mL	100 mg/5 mL	50 mg each	100 mg each	100 mg each	200 mg each		
	DOSE	DROPPER	TEASPOON OR SYRINGE	TABLET	TABLET	CAPLETS OR COATED TABLETS	TABLET OR CAPSULE		
WEIGHT (lbs)	AGE								
12-17	4-11 months	1.25 mL	1/2 tsp or 2.5 mL						
18-23	12-23 months	1.875 mL	3/4 tsp or 3.75 mL						
24-35	2-3 years		1 tsp or 5 mL	2 tablets					
36-47	4-5 years		1 1/2 tsp or 7.5 mL	3 tablets					
48-59	6-8 years		2 tsp or 10 mL	4 tablets	2 tablets	2 tablets			
60-71	9-10 years		2 1/2 tsp or 12.5 mL	5 tablets	2-2 1/2 tablets	2-2 1/2 tablets			
72-95	11 years		3 tsp or 15 mL	6 tablets	3 tablets	3 tablets	2 each		
over 96	12 years & older					4 tablets	2 each		